Amination Utilizing *N*-Haloimides as Nitrogen Source

Hongbo Li,^a Bao-Yi Ren,^{*,b} and Fu-Shun Liang^{*,a}

^a College of Chemistry, Liaoning University, Shenyang, Liaoning 110036, China

^b College of Science, Shenyang University of Chemical Technology Shenyang, Liaoning 110142, China

Email: renbaoyi@syuct.edu.cn (B.-Y. R.), fsliang@lnu.edu.cn (F.-S. L.)

Abstract Conventionally, *N*-bromosuccinimide (NBS) is a brominating and oxidizing reagent in organic synthesis. In our research on *N*-haloimides in organic transformation, we proposed the concept of utilizing *N*-haloimides like NBS as an amination reagent, *i.e.*, direct incorporation of imido unit to the target molecules to achieve amination, with the assistance of positive halogen(I). In this paper, we would like to summarize the amination of various fundamental substrates with *N*-haloimides as the amination reagents. We have developed an *N*-haloimide-DBU (Lewis base activator) combination system, which affords halogen bonding complex and subsequent ion pair intermediate *via* halogen transfer. All these reactions feature convenient and readily available reagents, mild conditions, simple execution, broad halogen scope, high efficiency and metal-free. The utilization of *N*-haloimides as versatile aminating reagents has been demonstrated and further application of the synthetic protocol for the preparation of bioactive and optical materials is expected.

Keywords amination, *N*-haloimides, halogen activation, nitrogen sources, halogen bond

Introduction

Research on new reagents, new reactions and new methods in synthetic chemistry is one of hot topics in modern chemistry. Hence, development of new reagents or exploration of new function of known reagents is of great significance.

From one hundred years ago to present, elemental bromine has been used to conduct bromination reaction. Later, *N*-bromophthalimide (NBP) was discovered as an efficient and easy-handled bromination reagent. In 1942, Ziegler utilized *N*-bromosuccinimide (NBS) to achieve bromination (Scheme 1).^[1] Afterwards, NBS has been introduced to the textbook of Organic Chemistry, as a typical and well-known brominating reagent for synthetic organic chemistry.

Scheme 1 Commonly used brominating reagents

Commercially available NBS can be regarded as convenient source of either a bromine radical used in radical reactions or cationic bromine used in electrophilic reactions (Scheme 2).^[2,3] In NBS, the halogen atom is attached to the nitrogen atom through a covalent bond with low electrophilicity and thus low reactivity. For the homolytic N—Br cleavage of NBS, light irradiation or heating to high temperature is necessary. While for the heterolytic N—Br cleavage, there are three types of halogen activation methods: Brønsted acid activation, Lewis acid activation and Lewis base (nucleophile) activation (Scheme 3). Both Brønsted acid and Lewis acid bind to oxygen atom of succinimido moiety, thereby enhancing the electrophilicity of bromine atom. For Lewis base activation, nucleophilic attacking of bromine atom directly brings about a highly positive bromine species.

Scheme 2 Cleavage patterns of NBS

Scheme 3 Halogen activation

Lewis base activation mode shown in Scheme 3 is closely related to halogen bonding and halogen bond complex. Halogen bonding, akin to the hydrogen bonding, is a noncovalent attractive interaction between an electrophilic region (σ -hole) on a halogen atom in a molecule (halogen bond donor) and a Lewis base functioning as the halogen bond acceptor (Scheme 4a).^[4] Scheme 4b gives some representative examples of halogen bond complexes.^[5] So far, the halogen bonding has been widely used in crystal engineering, materials science, molecular recognition and biological systems and drug design. However, the application of halogen bonding in synthetic organic chemistry has received relatively less attention.^[6]

Scheme 4 Halogen bonding and halogen bond complexes

a) Halogen Bonding

Y = C, N, halogen, etc.; X = I, Br, Cl
 D = N, O, S, Se, Cl, Br, I... or I⁺, Br⁺, Cl⁺, F⁺ etc.

b) Selected Examples of Halogen Bond Complexes

In halogen bond complexes, the intermolecular interaction is weak without significant charge transfer. There is another type of halogen bond complexes, in which the halogen bond acceptor (Lewis base) is very electron-donating. Thus, the interaction is so strong that it leads to complete charge separation. The halogen atom is transferrable and attached on the halogen acceptor. In this case, halogen atom is activated and can be used in further transformations such as halogenation. Selected examples of halogen activation by Lewis base are shown in Scheme 5. For example, NBS can be activated by triphenylphosphine or dimethylthioether to be the ion pair form.^[7a,7b] *N*-iodosaccharin can be activated by 4-dimethylaminopyridine (evidenced by single crystal X-ray structure).^[7c]

Scheme 5 Examples of halogen activation by Lewis base (Nu)

Proposal of *N*-Haloimide as Nitrogen Source

During the process of activating NBS for bromination, relatively inert succinimide would be liberated as a by-product. Atom economic utilization of bromine and/or succinimide of NBS would be highly desirable, but this type of reactivity mode is quite less documented. We questioned whether it might be possible to enable direct installment of the succinimido moiety of NBS into the target molecules, *i.e.*, to develop amination reactions, in which highly reactive bromo-species functions merely as an intermediate (Scheme 6).^[8] Literature search reveals that succinimidyl-, phthalimidyl- and sulfonamidyl moiety containing organic compounds or reagents have been reported as the nitrogen sources under certain reaction conditions (Scheme 7).^[9] Moreover, Sharpless group has pioneered the use of *N*-bromoacetamide as the nitrogen source in asymmetric aminohydroxylation early in 1997.^[10]

Scheme 6 Proposal of a new concept—Using NBS as an amination reagent

Inspired by the Corey-Kim oxidation,^[7a] we hypothesized to use halogen bonding and halogen activation strategy to achieve this goal, as exemplified in Scheme 8. The choice of

Scheme 7 Reported nitrogen sources with succinimidyl, phthalimidyl and sulfonamidyl moiety **Scheme 8** Establishment of the activation model with NBS/DBU combination

1,8-diazabicyclo[5.4.1]undec-7-ene (DBU) was based on the consideration of the double duty of DBU: both strong base and nucleophile.^[11] NBS (halogen donor) would react with DBU (halogen acceptor) to form a 1:1 halogen bond adduct *via* halophilic attacking.^[12] Due to the electron-donating ability of DBU and good nucleofuge property of succinimide, bromine transfer^[13] may occur, affording an ion pair intermediate **I** (resonance structure of aldimine in DBU is beneficial for the stabilization of the resulting ion pair). DBU-stabilized bromonium ion in intermediate **I**, is highly electrophilic^[14,15] and succinimide anion would be a potential *N*-nucleophile. In this way, DBU-activated NBS *via* halogen bond interaction brings about significantly enhanced electrophilic reactivity for bromine and nucleophilicity for an imido-nitrogen atom. This provides opportunity for the development of a body of conceptually new amination reactions, which rely on the employment of halogen(I) as an electrophilic center and succinimide as the source of subsequent nucleophilic amination.

In 1999, Legon forecasted halogen bond adduct is a “pre-reactive complex” in organic transformation, but lack of experimental support and deep mechanistic understanding.^[16] In 2017, Sekar *et al.* also pointed out that the halogen(I)-bonded adduct is a “pre-reactive complex” and not a thermodynamically stable intermediate.^[17]

In our experiment, a mixture of NBS and DBU in CH₂Cl₂ may form a highly polar complex, as observed on TLC plate, indicating the production of ion pair intermediate **I**. Compared with the ¹H NMR data of free NBS and DBU, the chemical shifts of the NBS/DBU mixture (equal equivalent) change to some extent (the protons on DBU move downfield and protons on NBS move toward upfield), indicating strong intermolecular interaction between NBS and DBU. The theoretical calculation result supports the proposed ion pair intermediate **I**, in which three molecules of water are included (Scheme 9).^[18] The role of the water is supposed to distribute the negative charge on the succinimide anion, thus stabilizing the ion pair formed. The ion-pair intermediate is also supported by measuring the ion

Scheme 9 DFT structure of ion-pair intermediate I

conductivities of NBS/DBU *versus* NBS in DMF. The ion conductivity for the former is two orders of magnitude higher than the latter.

Amination Using *N*-Haloimides as Nitrogen Source

Intrigued by the unique reactivity described in Scheme 8, we envisioned that it is possible to develop the *N*-haloimide/Lewis base combination to be one type of atom-efficient and versatile aminating reagent with the aid of halogen(I). With this idea in mind, we devoted our efforts to explore the reaction of various substrates with the *N*-haloimide/DBU system.

This section describes the amination by using *N*-haloimides as the nitrogen source with a broad spectrum of fundamental substrates including β -diketonate compounds, alkyl aryl ketones, α,β -unsaturated enones, alkenes, indoles, alkyl halides, P(H) compounds, nitrones, sulfones, and alkynes. Selected mechanism is also presented.

α -Amination of β -dicarbonyl compounds

α -Amido β -dicarbonyl compounds are versatile organic intermediates in the synthesis of various heterocyclic compounds, peptide mimetics, α -amino acids and their derivatives, and β -hydroxyl α -amino esters. Direct α -amination of the readily available β -dicarbonyl compounds is more convenient and highly desirable for the synthesis of α -amido β -dicarbonyl compounds. Thus, we aimed at the direct α -C—H imidation of β -dicarbonyl compounds by using DBU-activated *N*-haloimides as both electrophilic and nucleophilic reagents (Scheme 10). As a result, the *N*-haloimide/DBU combination provides a convenient and economic protocol toward α -C—H imidation of β -dicarbonyl compounds.^[19] One of the distinct features of the reaction is without the necessity of external nitrogen sources.

Under optimized conditions, we investigated this methodology with a range of β -dicarbonyl compounds (Scheme 10a). β -Keto esters, β -ketoamides, β -diketone and malonic esters all can afford the corresponding α -imidation products **2** in high yields. Products were isolated either in a keto/enol mixture or in complete enol/keto form. Similar to NBP, other *N*-haloimides like NBS and *N*-bromosaccharin were also competent reagents as potential self-immolating nitrogen sources (Schemes 10b and 10c). Thus, α -imidated compounds were obtained in moderate to high yields. Once again, keto/enol mixture or complete enol form was observed.

The possible mechanism for the α -imidation of β -dicarbonyl compounds was proposed (Scheme 11). Enolate **II** reacts with ion pair intermediate **I** to give α -bromo β -dicarbonyl **III**, followed by nucleophilic substitution by the phthalimide anion. The function of DBU is to deprotonate β -dicarbonyl compounds (as a base) and to activate NBP (as a nucleophilic promoter).

α -Amination of ketones

Generally, the procedure for α -amination of ketones requires multiple steps. In 2004, Lei *et al.* synthesized α -imido ketones in two steps (Scheme 12).^[20] In our study, a one-pot

Scheme 10 α -C—H amination of β -dicarbonyl compounds using DBU-activated *N*-haloimides **Scheme 11** Proposed mechanism for α -amination **Scheme 12** One-step *versus* two-step synthesis of α -amino ketones

cascade transformation of ketones into α -imido ketones has been developed by using NBS/DBU combination strategy.^[21] NBS provides both electrophilic bromine and nucleophilic nitrogen sources, and DBU functions as a base and a nucleophilic promoter for the activation of NBS. α -Bromination is considered to be the key step in the process, which takes

place between more electrophilic bromide active species and enolates.

A variety of ketones were reacted with NBS under optimized conditions (Scheme 13). Methyl aryl ketones containing both electron-donating groups and electron-withdrawing groups on the aryl rings could be smoothly transformed into the desired products **4**. 2-Naphthyl methyl ketone provided the corresponding product in high yield. Heteroaryl ketones, including furan, thiophene, and pyridine heterocycles, were converted into the target products in 81%–86% yields. 4-Phenylbut-3-en-2-one, propiophenone, butyrophenone and α -phenyl acetophenone were also suitable substrates, affording α -aminated products in high yields. Besides NBS, NBP was also a competent reagent, affording the desired isindole-1,3-diones in high yield. For the reaction of acetophenone with 1,3-dibromo-5,5-dimethylhydantoin, due to the reactivity of the bromine attached to the imine N-atom being higher than that on the amide N-atom, the corresponding α -imidated ketone was produced in 85% yield (debrominated).

Scheme 13 Direct α -amination of ketones by *N*-haloimides/DBU combination strategy

β -Amination of chalcones

In the continued work, a one-pot cascade transformation of chalcones into β -imidoketones has been developed.^[22] Under the optimal conditions, a range of reactions were carried out with **5** and NBS (1.2 equiv) in the presence of DBU (1.2 equiv) in MeCN (Scheme 14). The reactions proceeded smoothly to afford the corresponding β -imidoketones **6** in moderate to high yields. The R^2 group on the substrates **5** may be either electron-rich or electron-deficient aryl, heteroaryl and alkyl groups. The R^1 substituents were also broad, including electron-deficient aryl, electron-rich aryl, naphthyl and heteroaryl. The whole process involves tandem bromoamination/debromination, in which NBS functions as both electrophilic bromine and nitrogen sources and DBU as a nucleophilic promoter to activate NBS to be a more electrophilic bromine species and to remove the bromine of α -bromoketone. It was found that trace amount of water is crucial for the reaction to proceed.

Scheme 14 β -Amination of chalcones via a tandem bromoamination/debromination sequence

Later, other research groups have also achieved some interesting results. Lonea and Bhat reported a simple and straightforward method for the synthesis of a range of functionalized enamides from electron deficient alkenes using NBS and DABCO combination.^[23] Cai *et al.* developed an effective enantioselective β -amination of chalcones with NBS into β -imidoketones using a bis(oxazoline) chiral ligand.^[24]

Allylic amination of alkenes

Allylic amines represent an important structural motif frequently found in natural products, pharmaceuticals, as well as a versatile building block for the synthesis of organic molecules with a high degree of complexity. To date, commonly used approaches for the allylic amination of olefins include (i) metal-based oxidative allylic amination, and (ii) metal-free allylic amination, for example, by the introduction of a selenium reagent, or a hypervalent iodine(III) reagent (Scheme 15, top). We developed a novel and complementary access to allylic amination by the reaction of alkenes with an NBP or NBS/DBU combination under mild conditions (Scheme 15, bottom).^[25]

Scheme 15 Metal-based and metal-free allylic amination

Under optimized conditions, a range of reactions were carried out with various alkenes **7** and NBP (1.2 equiv) in the presence of DBU (1.2 equiv) in CH_2Cl_2 (Scheme 16). The alkene substrates investigated include substituted α -methylstyrenes (methyl, Cl and F), allylic benzene, 2,3-dimethylbutene, tetrasubstituted alkenes like tetramethylethene, cyclic alkenes such as 1-methylcyclohexene and 1-phenylcyclohexene. The corresponding allylamines **8** were obtained in moderate to excellent yields. External *N*-nucleophiles can also be introduced efficiently in the cases of the utilization of NBS/DBU combination. The scope of external amines appears broad, such as *O*-alkyl-*N*-arylsulfonylcarbamates (alkyls = methyl and benzyl; aryls = phenyl and 4-tolyl), *N*-methyltosylamine, phthalimide and substituted phthalimides (4-methyl and 4- NO_2), saccharin, 1*H*-benzo[d][1,2,3]triazole. It was found that the

Scheme 16 Allylic amination of alkenes using NBP(S)/DBU combination

Internal Nitrogen Source:

External Nitrogen Source:

installation of the external amino group is dependent on the pK_a values of the imides employed. For instance, succimide ($pK_a = 14.7$) is unable to be introduced, while amines having a more acidic NH group, can be used as the nitrogen sources in the allylic amination.

A possible mechanism for the allylic amination of alkenes was proposed (Scheme 17). Firstly, alkene reacts with ion pair **I** to furnish a DBU-stabilized bromonium ion **II** and its open form **III**. Secondly, allyl bromide **IV** is formed *via* proton elimination. Finally, nucleophilic substitution by the phthalimide anion leads to allylamine **8**.

Scheme 17 Proposed mechanism for the allylic amination

In 2016, Shinde and Kshirsagar reported a NBS promoted and base switchable reaction with styrene substrates and α -imido and α -amino ketones were obtained under elevated temperature.^[26]

α -Amination of indole derivatives

After the success of amination with alkene substrates, we further explored direct imidation of indole derivatives enabled by DBU-activated *N*-haloimides (Scheme 18).^[27]

Scheme 18 α -Amination of *N*-substituted indoles using *N*-haloimides as the nitrogen sources

N-methylindole substrates containing either electron-donating methyl, methoxy or weak electron-withdrawing groups like halogens (F, Cl, and Br) on the phenyl ring of the indole skeleton, afforded the corresponding imidated products **10** in moderate yields. As for strong electron-withdrawing groups like methoxycarbonyl and nitro- on the phenyl ring of *N*-methylindoles, the yields were fairly low. The corresponding 3-bromo-*N*-methylindole derivatives were proved to be the main products. In the case of a methyl group located at the 3-position of *N*-methylindole, imidated product was obtained in 91% yield. However, electron-withdrawing methoxycarbonyl group at the same position gave no reaction. As can be seen from above, electron-rich groups on the indole ring displayed higher reactivity than electron-withdrawing groups in this reaction. *N*-Benzyl and *N*-allylindoles afforded products in respective 56% and 72% yields. The allyl functional group exhibits a good tolerance. *N*-methylpyrrole was also efficient to afford the target product in 62% yield. The scope of the reaction was further investigated by exploring other types of *N*-haloimides as potential self-immolating nitrogen sources. The reaction of *N*-methylindole with *N*-bromophthalimide (2.2 equiv), or *N*-bromosaccharin in the presence of DBU (2.2 equiv) gave the imidated product at room temperature. All the reactions proceeded rapidly and completed within 10 min. The amination process involves tandem bromonium ion formation, electrophilic addition and elimination of HBr.

Amination of halides: DBU-mediated C—N coupling reaction

Amine is important raw material and intermediate in medicine, pesticide, as well as liquid crystal material. Phthalimides are versatile synthetic tools used mainly for the preparation of amines *via* the Gabriel reaction.^[28] In this line, we developed a new type of cross-coupling reaction between alkyl halides and *N*-haloimides, which provides an efficient and straightforward method for the synthesis of amines and functionalized amines.^[29] We found that, with organobases like DBU as the activator, benzyl chloride can react with *N*-chlorophthalimide

Scheme 19 Reaction of benzyl chloride with NCP **Scheme 20** Proposed mechanism for the C—N coupling of halides and *N*-haloimides

(NCP), giving the corresponding imidation product in 91% yield. No reaction would occur in the absence of DBU (Scheme 19). The proposed mechanism for the imidation involves the nucleophilic substitution of alkyl halides by phthalimide anion (Scheme 20). The formation of dihalogen byproduct was proved by KI-starch test paper experiment. We suggest that the halogen bond between intermediate II and RX may facilitate the nucleophilic substitution reaction. The mutual activation induced by halogen bonding makes otherwise inert reaction to take place.

The scope of this coupling reaction is fairly broad (Scheme 21). Alkyl chlorides examined include benzyl chloride, *n*-butyl chloride, α -chlorocyclopentanone and α -chloro *N,N*-dimethylacetamine. Benzyl chloride is highly efficient, while *n*-butyl chloride exhibits relatively lower efficiency. α -Chlorocyclopentanone and α -chloro *N,N*-dimethylacetamine gave satisfactory yields. Primary alkyl bromides containing various substituents like benzyl, allyl, ethyl, homobenzyl, isobutyl are suitable substrates. A lot of functional groups are compatible with the reaction conditions, such as methoxy, dimethoxyacetal, carbonyl and ester. Secondary alkyl bromides like isopropyl bromide and diphenylmethyl bromide can also afford the corresponding aminated product, albeit in low yields. However, cyclic alkyl bromide, *i.e.*, cyclohexyl bromide, was inefficient. Alkyl iodide like methyl iodide worked very well.

Dihalides are also suitable for the cross-coupling reaction (Scheme 22). In the reaction of 1,2-dibromoethane with NCP, 1.2 equiv of DBU gives monobromo substituted amine in 56% yield, while enamide product was produced in 53% yield, in the presence of 2.2 equiv of DBU. For 1,4-dibromobutane, 1,6-dibromooctane and 1,2-dichloroethane substrates, the corresponding imidation products with a pendant bromine or chlorine atom were attained, which may be potentially useful in further transformation. Mixed dihalides were also subjected to the reaction sequence. For example, 1-bromo-4-chlorobutane gave chlorine atom-tolerant imidation product as the only product, due to the different reactivity between bromine and chlorine atoms.

Different *N*-haloimides exhibit different reactivity. NBP is less efficient than NCP. Similar trend was observed for *N*-halosuccinimides. *N*-Bromosaccharin afford 2-benzylbenzo-[*d*]isothiazol-3(2*H*)-one 1,1-dioxide in 89% yield.

Amination of phosphonates: DBU-mediated P—N cross-coupling

As an extension of our interest in developing new reactions

Scheme 21 DBU-mediated cross-coupling reaction of alkyl halides and NCP **Scheme 22** Cross-coupling reaction of dihalides with NCP

using *N*-haloimide as the nitrogen source, metal-free P—N cross-coupling of P(O)—H compounds with NBP was developed.^[30] DBU mediated coupling reaction between ethyl phenylphosphinate and NBP in MeCN at 0 °C has been proved to be the most efficient, furnishing the target product in 85% yield within 10 min.

The scope of the P(O)—H components in the P—N coupling reaction is broad and the reaction time is quite short (less than 10 min) (Scheme 23). Trivalent phosphorous substrates include *H*-phosphinates (both cyclic and acyclic), *H*-phosphonates with alkoxy, CF₃-containing alkoxy and aryloxy substituents on the P center, and diarylphosphine oxides. Phosphoramidate products **14** were obtained in moderate to high yields, which are of biological importance.

In a control experiment, TEMPO (1.5 equiv) was added immediately to a mixture of ethyl phenylphosphinate, NBP and DBU in MeCN at 0 °C (Scheme 24). Compared to the reaction without TEMPO, the yield has almost no change. The result may help one to exclude the possible radical pathway. Thus, electrophilic bromination of P(O)—H compounds gives rise to phosphoryl bromide II. Further addition/elimination affords the corresponding cross-coupled product.

Scheme 23 P—N Cross-coupling: Scope of P(O)—H

substrates

Scheme 24 Proposed mechanism for the P—N cross-coupling

a) Mechanistic study

b) Proposed mechanism

α -Amination of nitrones

In nitron chemistry, the synthesis of structurally interesting new derivatives have attracted considerable attention. Nitrones bearing a carbo- or heteroatom adjacent to the nitron group carbon atom are prepared either from nitron substrates *via* direct transformation or non-nitron precursors through indirect pathways, while metal-free methods toward direct α -functionalization of nitrones are less-documented.

DABCO mediated reactions of nitrones and NBS have been reported by Lobo *et al.* (Scheme 25, top).^[31] The reaction generally takes 2 d to finish, and the yields are fairly moderate (40%–64%). By contrast, DBU mediated direct α -amination of

Scheme 25 Direct α -amination of nitrones with the NBS(P)-organobase protocol

a) Lobo *et al.*

b) Our work

Scheme 26 Scope of nitrones and *N*-haloimides

nitrones has been developed (Scheme 25, bottom), which exhibited a broader scope (both for nitron and quinoline *N*-oxide substrates and *N*-haloimides), shorter reaction time (less than 5 h), and higher yields (62%–93%, Scheme 26).^[32]

A possible mechanism for α -amination of nitrones was proposed (Scheme 27). Nitron would first react with the highly electrophilic bromine species **I** to furnish an iminium ion **II**. Then, nucleophilic attack by the phthalimide anion gives rise to intermediate **III**. Finally, α -aminonitron **16** is produced *via* the elimination of DBU·HBr.

Scheme 27 Mechanistic proposal for α -amination of nitrones

Reductive amination of dimethyl sulfoxide

This section mainly explores the activation of dimethyl sulfoxide (DMSO) by halogen bond. In this case, Lewis bases like DBU was unnecessary, because of the internal halogen bonding interaction and halogen activation between *N*-haloimide and DMSO. Two examples of reductive amination were given in Scheme 28, in which NBP and *N*-bromosaccharin can directly react with DMSO at 80 °C, affording *N,S*-acetyls **17a** and **17b** in 72% and 38% yields, respectively. The reaction was supposed to undergo an initial ion pair intermediate **I**,^[33] subsequent formation of sulfonium **II** and final capture of which by *N*-nucleophile.

Scheme 28 Reductive amination of dimethyl sulfoxide

In 2016, Sudalai group reported the efficient transformation of alkenes and enol ethers into α -amino ketones and esters by applying DMSO-activated NBS.^[34]

Reaction with alkynes

The reaction of alkynes with *N*-haloimides is summarized in Scheme 29. Directly from terminal alkynes, 1-haloalkynes were prepared using DBU activated *N*-haloimides at room temperature (Scheme 29, top left). Under elevated temperature, the difunctionalization of terminal alkynes takes place, giving rise to useful halogenated enamines (Scheme 29, top right). Comparatively, utilizing DMF-activated *N*-haloimides, another type of difunctionalization product (*i.e.*, α -amino ketones) can be attained, simply by varying the feed sequence of the above-mentioned substrates (Scheme 29, bottom).

Scheme 29 Influence of different activation mode on the outcome of alkynes

Preparation of 1-haloalkynes

Directly from terminal alkynes and with *N*-halosuccinimides (halo = Br and I) or *N*-chlorophthalimide as the halogen sources, DBU as the activator, 1-haloalkynes **19** were prepared in good to excellent yields at room temperature (Scheme 30).^[35] Bis(bromoalkyne) and bis(iodoalkyne) were also synthesized in excellent yields with the NBS(NIS)/DBU combination possessing broad halogen scope. Compared to the literature reported methods, this synthetic strategy provided a greener and practical alternative towards 1-haloalkynes with simple execution, extremely short reaction time, broad halogen scope, high efficiency and metal-free.

Scheme 30 Halogenation of terminal alkynes

Alkyne aminohalogenation

If the above-mentioned reaction of terminal alkynes, DBU, and *N*-haloimides were carried out at elevated temperature, halogenated enamines **20** would be generated in high efficiency and high regio- and stereoselectivities (Scheme 31).^[36] This aminohalogenation process provided a new idea for the regio- and stereoselective synthesis of *cis*-halogenated enamines. The possible mechanism involves the initial formation of alkynyl

Scheme 31 Scope of alkynes and *N*-haloimides

Scheme 32 Mechanistic proposal for alkyne amino-halogenation

halides **19**, subsequent nucleophilic addition and final protonation (Scheme 32).

Alkyne aminooxygenation

Using DMF to activate NBS and adding DBU (as the base only) in the second step gives rise to a one-pot, two-step cascade reaction for the synthesis of α -amino ketones.^[37] Both terminal and internal alkynes are active, giving imidated ketones **22** efficiently. The *N*-source arises from inherent *N*-haloimides and the *O*-source is from water in the solvent. Monoamino ketones **23/24** were the main products in the reaction with NBS (Scheme 33). Mono and/or diamino ketones are produced when NBP is utilized (Scheme 34). A mechanism of sequential oxy-1,1-dibromogeneration of alkynes, nucleophilic substitution and reductive debromination (or second nucleophilic substitution) is proposed.

Nature of Amination

Above we have demonstrated that, direct installation of nitrogen functionality of *N*-haloimides into the target molecules is feasible using the *N*-haloimide–DBU combination strategy. The nature for the amination can be elucidated as the following two aspects (Scheme 35). On one hand, in the case of *R*–H type of substrates, halogenation takes place first, followed by

Scheme 33 Aminohydroxylation of alkynes *via* DMF-activated *N*-haloimides. **Scheme 34** Reactions of alkynes with NBP **Scheme 35** Two ways involved in the amination reactions

nucleophilic substitution (pathway 1). On the other hand, for alkene-, alkyne-, indole-type substrates, haloiminium intermediate would be produced first. Subsequent nucleophilic attacking as well as elimination leads to the aminated products (pathway 2).

In the above amination reactions, DBU-mediated reactions between halides/aryl alkyl ketones and *N*-haloimides involve a nice combination of two known name reactions: Corey-Kim oxidation (initial activation step)^[7a] and Gabriel synthesis (Scheme 36).^[28]

Scheme 36 Comparison of the reaction developed here with known name reactions

a) Corey-Kim oxidation (activation step):

b) Gabriel Synthesis:

c) Reactions described in this Account

In the amination reaction, we explored the scope of each component within *N*-haloimide-Lewis base system. Appropriate *N*-haloimide-based nitrogen sources include *N*-halosuccinimides (NBS, NIS, NCS), *N*-bromophthalimide (NBP), NCP, 1,3-dibromo-5,5-dimethylhydantoin and *N*-bromosaccharin, and so on (Scheme 37a). In addition to using DBU as an efficient halogen activator and moderate to strong base, 1,5-diazabicyclo[4.3.0]non-5-ene (DBN), DMF and DMSO, etc., can also be used to activate *N*-haloimides (Scheme 37b). For different type of substrates, *N*-haloimide-Lewis base combination displays slightly different reactivity. The success of incorporating *N*-nucleophile in the target molecules depends on the nucleofuge ability and/or nucleophilicity of *N*-haloimides.

Scheme 37 Summary of (a) the scope of the nitrogen source regarding *N*-haloimides and analogs. (b) Scope of Lewis bases (activators)

Conclusions and Perspectives

Herein, we summarize our efforts on the development of amination reactions utilizing *N*-haloimides as the nitrogen source. We have developed a broad spectrum of fundamental substrates including β -diketone compounds, alkyl aryl ketones, α,β -unsaturated enones, alkenes, *N*-substituted indoles, alkyl halides, P(H) compounds, nitrones, dimethylsulfone, and alkynes. All these metal-free reactions feature convenient and readily available reagents, mild conditions, simple execution, high efficiency and atom-economy.

We disclosed that Lewis base-activated *N*-haloimides can be utilized as an efficient amination reagent. Appropriate *N*-haloimide-based nitrogen sources include *N*-halosuccinimides (NBS, NIS, NCS), NBP, NCP, 1,3-dibromo-5,5-dimethylhydantoin and *N*-bromosaccharin. Similar to hydrogen bond activation, the halogen bonding and halogen

activation was highlighted herein. For the nature of amination, two main ways can be distinguished. In the case of R—H type substrates, halogenation takes place first, followed by nucleophilic substitution; For alkene-, alkyne-, indole-type substrates, haloiminium intermediate would be produced first. Subsequent ring-opening as well as elimination lead to aminated products. Considering the amination reactions were mostly conducted at room temperature without light irradiation, along with the control experiments using radical scavenger, halogen activated ionic mechanism, rather than radical pathway, would be the most probable.

The work reported in this paper supports our proposed concept of amination reactions utilizing *N*-haloimides as the nitrogen source, rather than traditional concept of *N*-haloimides as the halogenating reagents. The significance of the study would be, on one hand, open new viewpoints of developing new function of known reagents for new reactions. On the other hand, it may find potential applications in the life science fields, because of the importance of amination reaction and imido moieties in pharmaceuticals and agrochemicals.

As for the limitation and extension of the substrate scope, it is noteworthy that the amination of arenes and heteroarenes cannot be achieved using this strategy, because the aryl-halogen bond in the halogenation product is inert and the S_NAr reaction cannot occur. Interestingly, visible light-induced C—H imination of aryl groups with NCP has been achieved by nitrogen-centered radical-mediated reactions of aryl groups with NCP at room temperature.^[38]

The halogen activation mode proposed in this paper can be adopted to explain the reactivity of relevant reactions. It may also help to design and perform useful organic reactions. Lin and John have discovered an oxidative homo-coupling method for the synthesis of symmetric azobenzenes from substituted anilines, employing NCS and DBU reagents.^[39] Wang group has reported an aminotrifluoromethylthiolation of α,β -unsaturated carbonyl compounds with DABCO/*N*-trifluoromethylthio-4-nitrophthalimide combination.^[40]

For the future, one can expect more unprecedented reactions to be developed based on the exciting *N*-haloimide/ Lewis base combination system, either in the absence or presence of external nucleophiles. The application of this synthetic protocols for the preparation of various bioactive or optical compounds is also worth awaiting.

Acknowledgement

Financial support from the Liaoning Revitalization Talents Program (No. XLYC1902111), the Key projects of Department of Education, Liaoning Province (No. LZD202005), the Science Research Plan of SYUCT (No. XXLJ2019006), and the Natural Science Foundation of Liaoning Province (No. 2021-MS-254) is gratefully acknowledged.

Author Contributions

H. L. and B.-Y. R. wrote this manuscript. F.-S. L. initiated the project and polished this manuscript.

Conflict of Interest

The authors declare no conflict of interest.

Copyright © 2021 Hongbo Li, Bao-Yi Ren, Fu-Shun Liang. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<http://creativecommons.org/licenses/by/4.0/>). The use, distribution or reproduction in other forums is permitted, provided the original author(s) or licensor are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or

reproduction is permitted which does not comply with these terms.

References

- [1] (a) Wohl, A. Bromierung ungesättigter Verbindungen mit *N*-Brom-acetamid, ein Beitrag zur Lehre vom Verlauf chemischer Vorgänge, *Ber. Dtsch. Chem. Ges. B* **1919**, 52, 51–63; (b) Ziegler, K.; Späth, A.; Schf, E.; Schumann, W.; Winkelmann, E. *Die Halogenierung ungesättigter Substanzen in der Allylstellung. Justus Liebigs Annalen Der Chemie* **1942**, 551, 80–119; (c) Djerassi, C. Brominations with *N*-Bromosuccinimide and Related Compounds. The Wohl-Ziegler Reaction. *Chem. Rev.* **1948**, 43, 271–317; (d) Pearson, R. E.; Martin, J. C. The Identity of the Chain-Carrying Species in Brominations with *N*-Bromosuccinimide: Selectivity of Substituted *N*-Bromosuccinimides toward Substituted Toluenes. *J. Am. Chem. Soc.* **1963**, 85, 3142–3146; (e) Koval, I. V. *N*-Halo Reagents. *N*-halo reagents. *N*-halosuccinimides in organic synthesis and in chemistry of natural compounds. *Russ. J. Org. Chem.* **2002**, 38, 301–337.
- [2] (a) Sasaki, M.; Yudin, A. K. Oxidative Cycloamination of Olefins with Aziridines as a Versatile Route to Saturated Nitrogen-Containing Heterocycles. *J. Am. Chem. Soc.* **2003**, 125, 14242–14243; (b) Zhou, L.; Tan, C. K.; Zhou, J.; Yeung, Y.-Y. Ling Zhou. Facile, Efficient, and Catalyst-Free Electrophilic Aminoalkoxylation of Olefins: Scope and Application. *J. Am. Chem. Soc.* **2010**, 132, 10245–10247; (d) Zhou, L.; Tan, C. K.; Jiang, X.; Chen, F.; Yeung, Y.-Y. Asymmetric Bromolactonization Using Amino-thiocarbamate Catalyst. *J. Am. Chem. Soc.* **2010**, 132, 15474–15476; (e) Cai, Y.; Liu, X.; Hui, Y.; Jiang, J.; Wang, W.; Chen, W.; Lin, L.; Feng, X. Catalytic Asymmetric Bromoamination of Chalcones: Highly Efficient Synthesis of Chiral α -Bromo- β -Amino Ketone Derivatives. *Angew. Chem., Int. Ed.* **2010**, 49, 6160–6164.
- [3] (a) Horner, L.; Winkelmann, E. H. Neuere Methoden der präparativen organischen Chemie II 14. *N*-Bromsuccinimid, Eigenschaften und Reaktionsweisen Studien zum Ablauf der Substitution XV. *Angew. Chem.* **1959**, 71, 349–365; (b) Walling, C.; Rieger, A. L.; Tanner, D. D. Positive Halogen Compounds. VIII. Structure and Reactivity in *N*-Bromosuccinimide Brominations. *J. Am. Chem. Soc.* **1963**, 85, 3129–3134; (c) Russell, G. A.; Desmond, K. M. Directive Effects in Aliphatic Substitutions. XIX. Photobromination with *N*-Bromosuccinimide. *J. Am. Chem. Soc.* **1963**, 85, 3139–3141.
- [4] (a) Eds.: Metrangolo, P.; Resnati, G. *Halogen Bonding: Fundamentals and Applications*. Springer, Berlin, **2008**; (b) Eds.: Metrangolo, P.; Resnati, G. *Halogen Bonding vol I and vol II. Impact on Materials Chemistry and Life Sciences, Topics in Current Chemistry Series 358*, Springer, Switzerland, **2015**; (c) Cavallo, G.; Metrangolo, P.; Milani, R.; Pilati, T.; Priimagi, A.; Resnati, G.; Terraneo, G. The Halogen Bond. *Chem. Rev.* **2016**, 116, 2478–2601; (d) Bulfield, D.; Huber, S. M. Halogen Bonding in Organic Synthesis and Organocatalysis. *Chem.-Eur. J.* **2016**, 22, 14434–14450; (e) Wang, H.; Wang, W.; Jin, W. J. σ -Hole bond vs. π -hole bond: a comparison based on halogen bond. *Chem. Rev.* **2016**, 116, 5072–5104.
- [5] (a) Castellote, I.; Morón, M.; Burgos, C.; Alvarez-Builla, J.; Martin, A.; Gómez-Sal, P.; Vaquero, J. J. Reaction of imines with *N*-iodosuccinimide (NIS): unexpected formation of stable 1 : 1 complexes. *Chem. Commun.* **2007**, 1281–1283; (b) Crowston, E. H.; Lobo, A. M.; Prabhakar, S.; Rzepa, H. S.; Williams, D. J. X-Ray and SCF-MO model study of the complex formed between *N*-bromosuccinimide and 1,4-diazabicyclo[2.2.2]octane. *Chem. Commun.* **1984**, 276–278; (c) Raatikainen, K.; Rissanen, K. Breathing molecular crystals: halogen- and hydrogen-bonded porous molecular crystals with solvent induced adaptation of the nanosized channels. *Chem. Sci.* **2012**, 3, 1235–1239; (d) Wang, Y.-M.; Wu, J.; Hoong, C.; Rauniyar, V.; Toste, F. D. Enantioselective

- halocyclization using reagents tailored for chiral anion phase-transfer catalysis. *J. Am. Chem. Soc.* **2012**, *134*, 12928–12931; (e) Sladojevich, F.; McNeill, E.; Börgel, J.; Zheng, S.-L.; Ritter, T. Condensed-Phase, Halogen-Bonded CF_3I and $\text{C}_2\text{F}_5\text{I}$ Adducts for Perfluoroalkylation Reactions. *Angew. Chem. Int. Ed.*, **2015**, *54*, 3712–3716; (f) Tan, C. K.; Zhou, L.; Yeung, Y.-Y. Condensed-Phase, Halogen-Bonded CF_3I and $\text{C}_2\text{F}_5\text{I}$ Adducts for Perfluoroalkylation Reactions. *Synlett* **2011**, 1335–1339; (g) X. Sun, W. Wang, Y. Li, J. Ma, S. Yu, Halogen-bond-promoted double radical isocyanide insertion under visible-light irradiation: synthesis of 2-fluoroalkylated quinoxalines. *Org. Lett.* **2016**, *18*, 4638–4641.
- [6] (a) Bruckmann, A.; Pena, M. A.; Bolm, C. Organocatalysis through Halogen-Bond Activation. *Synlett* **2008**, 6, 900–902; (b) Coulem-bier, O.; Meyer, F.; Dubois, P. Controlled room temperature ROP of *L*-lactide by ICl_3 : a simple halogen-bonding catalyst. *Polym. Chem.*, **2010**, *1*, 434–437; (c) Walter, S. M.; Kniep, F.; Herdtweck, E.; Huber, S. M. Halogen-Bond-Induced Activation of a Carbon-Heteroatom Bond. *Angew. Chem., Int. Ed.* **2011**, *50*, 7187–7191; (d) Jungbauer, S. H.; Schindler, S.; Kniep, F.; Walter, S. M.; Rout L.; Huber, S. M. Multidentate halogen-bond donors as Lewis acidic activators or catalysts in halide abstraction reactions. *Synlett* **2013**, 24, 2624–2628.
- [7] (a) Corey, E. J.; Kim, C. U. New and highly effective method for the oxidation of primary and secondary alcohols to carbonyl compounds. *J. Am. Chem. Soc.* **1972**, *94*, 7586–7587; (b) Sakakura, A.; Ukai, A.; Ishihara, K. Enantioselective halocyclization of poly-prenoids induced by nucleophilic phosphoramidites. *Nature* **2007**, *445*, 900–903; (c) Makhotkina, O.; Liefbrig, J.; Jeannin, O.; Four-migué, M.; Aubert, E.; Espinosa, E. Cocrystal or salt: solid state-controlled iodine shift in crystalline halogen-bonded systems. *Cryst. Growth Des.* **2015**, *15*, 3464–3473.
- [8] (a) Day, J. C.; Katsaros, M. G.; Kocher, W. D.; Scott A. E.; Skell, P. S. Addition reactions of imidyl radicals with olefins and arenes. *J. Am. Chem. Soc.* **1978**, *100*, 1950–1951; (b) Alix, A.; Lalli, C.; Re-tailleau, P.; Masson, G. Highly enantioselective electrophilic α -bromination of enecarbamates: chiral phosphoric acid and calcium phosphate salt catalysts. *J. Am. Chem. Soc.* **2012**, *134*, 10389–10392.
- [9] (a) Foo, K.; Sella, E.; Thomé, I.; Eastgate, M. D.; Baran, P. S. A mild ferrocene-catalyzed C-H imidation of (hetero)arenes. *J. Am. Chem. Soc.* **2014**, *136*, 5279–5282; (b) Lardy, S. W.; Schmidt, V. A. Intermolecular Radical Mediated Anti-Markovnikov Alkene Hydroamination Using *N*-Hydroxyphthalimide. *J. Am. Chem. Soc.* **2018**, *140*, 12318–12322; (c) Souto, J. A.; Zian, D. Muñiz, K. Iodine (III)-mediated intermolecular allylic amination under metal-free conditions. *J. Am. Chem. Soc.* **2012**, *134*, 7242–7245; (d) Li, Y.; Zhang, Q. *N*-Fluorobenzenesulfonimide: an efficient nitrogen source for C–N bond formation. *Synthesis* **2015**, 47, 159–174; (e) Muñiz, K. Promoting intermolecular C–N bond formation under the auspices of iodine (III). *Acc. Chem. Res.* **2018**, *51*, 1507–1519.
- [10] Bruncko, M.; Schlingloff, G.; Sharpless, K. B. *N*-Bromoacetamide—A New Nitrogen Source for the Catalytic Asymmetric Aminohydroxylation of Olefins. *Angew. Chem. Int. Ed.* **1997**, *36*, 1483–1486.
- [11] (a) Reed, R.; Reau, R.; Dahan, F.; Bertrand, G. DBU and DBN are Strong Nucleophiles: X-Ray Crystal Structures of Onio- and Dioxio-Substituted Phosphanes. *Angew. Chem., Int. Ed.* **1993**, 399–401; (b) A. K. Morri, Y. Thummala, V. R. Doddi, The dual role of 1, 8-diazabicyclo[5.4.0] undec-7-ene (DBU) in the synthesis of terminal aryl- and styryl-acetylenes via Umpolung reactivity. *Org. Lett.* **2015**, *17*, 4640–4643; (c) S. K. Moodapelly, G. V. M. Sharma, V. R. Doddi. Controlled Reactivity of 1,8-Diazabicyclo [5.4.0] undec-7-ene (DBU) in the Selective Synthesis of 1-(Bromoethynyl) arenes. *Adv. Synth. Catal.* **2017**, *359*, 1535–1540.
- [12] (a) Zefirov, N. S.; Makhon'kov, D. I. X-philic reactions. *Chem. Rev.*, **1982**, *82*, 615–624; (b) Grinblat, J.; Ben-Zion, M.; Hoz, S. Halophilic reactions: anomalies in bromine transfer reactions. *J. Am. Chem. Soc.* **2001**, *123*, 10738–10739.
- [13] (a) Brown, R. S. Investigation of the early steps in electrophilic bromination through the study of the reaction with sterically encumbered olefins. *Acc. Chem. Res.* **1997**, *30*, 131–137; (b) Cui, X.-L.; Brown, R. S. Mechanistic evaluation of the halocyclization of 4-penten-1-ol by some bis (2-substituted pyridine) and bis (2, 6-disubstituted pyridine) bromonium triflates. *J. Org. Chem.* **2000**, *65*, 5653–5658.
- [14] Gao, W.-C.; Hu, F.; Huo, Y.-M.; Chang, H.-H.; Li, X.; Wei, W.-L. I_2 -Catalyzed C—O Bond Formation and Dehydrogenation: Facile Synthesis of Oxazolines and Oxazoles Controlled by Bases. *Org. Lett.* **2015**, *17*, 3914–3917.
- [15] (a) Denmark, S. E.; Burk, M. T. Lewis base catalysis of bromo- and iodolactonization, and cycloetherification. *Proc. Nat. Acad. Sci. U. S. A.* **2010**, *107*, 20655–20660; (b) Denmark, S. E.; Kuester, W. E.; Burk, M. T. Catalytic, asymmetric halofunctionalization of alkenes—a critical perspective. *Angew. Chem., Int. Ed.* **2012**, *51*, 10938–10953; (c) Cheng, Y. A.; Chen, T.; Tan, C. K.; Heng, J. J.; Yeung, Y.-Y. Efficient medium ring size bromolactonization using a sulfur-based zwitterionic organocatalyst. *J. Am. Chem. Soc.* **2012**, *134*, 16492–16495.
- [16] Legon, A. C. Prereactive complexes of dihalogens XY with Lewis bases B in the gas phase: a systematic case for the halogen analogue $\text{B} \cdots \text{XY}$ of the hydrogen bond $\text{B} \cdots \text{HX}$. *Angew. Chem., Int. Ed.* **1999**, *38*, 2686–2714.
- [17] Guha, S.; Kazi, I.; Nandy, A.; Sekar, G. Role of Lewis-Base-Coordinated Halogen (II) Intermediates in Organic Synthesis: The Journey from Unstable Intermediates to Versatile Reagents. *Eur. J. Org. Chem.* **2017**, 5497–5518.
- [18] Yuan H.; Zhang, J. Mechanistic insights on DBU catalyzed β -amination of nbs to chalcone driving by water: Multiple roles of water. *J. Comput. Chem.* **2017**, *38*, 438–445.
- [19] Tan, H.; Li, M.; Liang, F. Direct α -CH amination of β -dicarbonyl compounds using DBU-activated *N*-haloimides as nitrogen sources. *RSC Adv.* **2014**, *4*, 33765–33768.
- [20] Lei, A.; Wu, S.; He, M.; Zhang, X. Highly Enantioselective asymmetric hydrogenation of α -phthalimide ketone: an efficient entry to enantiomerically pure amino alcohols. *J. Am. Chem. Soc.* **2004**, *126*, 1626–1627.
- [21] Wei, Y.; Lin, S.; Liang, F. One-Pot Cascade Leading to Direct α -Imidation of Ketones by a Combination of *N*-Bromosuccinimide and 1,8-Diazabicyclo[5.4.1]undec-7-ene. *Org. Lett.* **2012**, *14*, 4202–4205.
- [22] Wei, Y.; Lin, S.; Liang, F.; Zhang, J. *N*-Bromosuccinimide/1,8-Diazabicyclo[5.4.1]undec-7-ene Combination: β -Amination of Chalcones via a Tandem Bromoamination/Debromination Sequence. *Org. Lett.* **2013**, *15*, 852–855.
- [23] Lonea, A. M.; Bhat, B. Metal free stereoselective synthesis of functionalized enamides. *Org. Biomol. Chem.* **2014**, *12*, 242–246.
- [24] Deng, T.; Wang, H.; Cai, C. Application of bis(oxazoline) in asymmetric β -amination of chalcones. *New J. Chem.* **2015**, *39*, 102–105.
- [25] Wei, Y.; Liang, F.; Zhang, X. *N*-Bromoimide/DBU Combination as a New Strategy for Intermolecular Allylic Amination. *Org. Lett.* **2013**, *15*, 5186–5189.
- [26] Shinde, M. H.; Kshirsagar, U. A. *N*-Bromosuccinimide promoted and base switchable one pot synthesis of α -imido and α -amino ketones from styrenes. *Org. Biomol. Chem.* **2016**, *14*, 858–861.
- [27] Li, Y.; Zhang, L.; Yuan, H.; Liang, F.; Zhang, J. Rapid α -Amination of *N*-Substituted Indoles by Using DBU-Activated *N*-Haloimides as Nitrogen Sources. *Synlett* **2015**, 116–122.
- [28] (a) Gibson, M. S.; Bradshaw, R. W. The Gabriel synthesis of primary amines. *Angew. Chem. Int. Ed.* **1968**, *7*, 919–930; (b) Sheehan, J. C.; Bolhofer, W. A. An improved procedure for the condensation of potassium phthalimide with organic halides. *J. Am. Chem. Soc.* **1950**, *72*, 2786–2788.
- [29] Zhang, L.; Li, Y.; Jin, L.-Y.; Liang, F. Metal-free C–N cross-coupling of electrophilic compounds and *N*-haloimides. *RSC Adv.* **2015**, *5*,

- 65600–65603.
- [30] Li, Y.; Liang, F. With DBU-activated *N*-bromophthalimide as potential N-sources to achieve P—N cross-coupling of P(O)—H compounds. *Tetrahedron Lett.* **2016**, *57*, 2931–2934.
- [31] Lobo, A. M.; Prabhakar, S.; Rzepa, H. S.; Skapski, A. C.; Tavers, M. R.; Widdowson, D. A. C-substitution reactions of *c*, *n*-diaryl nitrones. *Tetrahedron* **1983**, *39*, 3833–3841.
- [32] Li, M.; Liang, F. Direct α -amination of nitrones achieved by DBU-activated *N*-haloimides. *Tetrahedron Lett.* **2016**, *57*, 3823–3826.
- [33] (a) Ashikari, Y.; Shimizu, A.; Nokami, T.; Yoshida, J. Halogen and chalcogen cation pools stabilized by DMSO. Versatile reagents for alkene difunctionalization. *J. Am. Chem. Soc.* **2013**, *135*, 16070–16073; (b) Shimizu, A.; Hayashi, R.; Ashikari, Y.; Nokami, T.; Yoshida, J. Switching the reaction pathways of electrochemically generated β -haloalkoxysulfonium ions-synthesis of halohydrins and epoxides. *Beilstein J. Org. Chem.* **2015**, *11*, 242–248; (c) Song, S.; Sun, X.; Li, X.; Yuan, Y.; Jiao, N. Efficient and Practical Oxidative Bromination and Iodination of Arenes and Heteroarenes with DMSO and Hydrogen Halide: A Mild Protocol for Late-Stage Functionalization. *Org. Lett.* **2015**, *17*, 2886–2889.
- [34] Prasad, P. K.; Reddi, R. N.; Sudalai, A. Regioselective Oxo-Amination of Alkenes and Enol Ethers with *N*-Bromosuccinimide-Dimethyl Sulfoxide Combination: A Facile Synthesis of α -Amino-Ketones and Esters. *Org. Lett.* **2016**, *18*, 500–503.
- [35] Li, M.; Li, Y.; Zhao, B.; Liang, F.; Jin, L.-Y. Facile and efficient synthesis of 1-haloalkynes via DBU-mediated reaction of terminal alkynes and *N*-haloimides under mild conditions. *RSC Adv.* **2014**, *4*, 30046–30049.
- [36] Li, M.; Yuan, H.; Zhao, B.; Liang, F.; Zhang, J. Alkyne aminohalogenation enabled by DBU-activated *N*-haloimides: direct synthesis of halogenated enamines. *Chem. Commun.* **2014**, *50*, 2360–2363.
- [37] Li, M.; Liang, F. One-pot, two-step conversion of alkynes to α -amino (α,α -diamino) ketones with a DMF-activated *N*-bromoimide strategy. *RSC Adv.* **2016**, *6*, 93325–93329.
- [38] Kim, H.; Kim, T.; Lee, D. G.; Roh, S. W.; Lee, C. Nitrogen-centered radical-mediated C-H imidation of arenes and heteroarenes via visible light induced photocatalysis. *Chem. Commun.* **2014**, *50*, 9273–9276.
- [39] John, A. A.; Lin, Q. Synthesis of Azobenzenes Using *N*-Chlorosuccinimide and 1,8-Diazabicyclo[5.4.0]undec-7-ene (DBU). *J. Org. Chem.* **2017**, *82*, 9873–9876.
- [40] Xiao, Q.; He, Q.; Li, J.; Wang, J. 1,4-Diazabicyclo[2.2.2]octane-Promoted Aminotrifluoromethylthiolation of α,β -Unsaturated Carbonyl Compounds: *N*-Trifluoromethylthio-4-nitrophthalimide Acts as Both the Nitrogen and SCF₃ Sources. *Org. Lett.* **2015**, *17*, 6090–6093.

Received November 4, 2021

Accepted November 15, 2021